



PLACE  
PICTURE  
HERE

Name: \_\_\_\_\_ D.O.B.: \_\_\_\_\_

Allergy to: \_\_\_\_\_

Weight: \_\_\_\_\_ lbs. Asthma: [ ] Yes (higher risk for a severe reaction) [ ] No

NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.

Extremely reactive to the following foods: \_\_\_\_\_

THEREFORE:

- [ ] If checked, give epinephrine immediately for ANY symptoms if the allergen was likely eaten.
- [ ] If checked, give epinephrine immediately if the allergen was definitely eaten, even if no symptoms are noted.

FOR ANY OF THE FOLLOWING:  
**SEVERE SYMPTOMS**



**LUNG**  
Short of breath,  
wheezing,  
repetitive cough



**HEART**  
Pale, blue,  
faint, weak  
pulse, dizzy



**THROAT**  
Tight, hoarse,  
trouble  
breathing/  
swallowing



**MOUTH**  
Significant  
swelling of the  
tongue and/or lips



**SKIN**  
Many hives over  
body, widespread  
redness



**GUT**  
Repetitive  
vomiting, severe  
diarrhea



**OTHER**  
Feeling  
something bad is  
about to happen,  
anxiety, confusion

OR A  
COMBINATION  
of symptoms  
from different  
body areas.



1. **INJECT EPINEPHRINE IMMEDIATELY.**
2. **Call 911.** Tell them the child is having anaphylaxis and may need epinephrine when they arrive.
  - Consider giving additional medications following epinephrine:
    - » Antihistamine
    - » Inhaler (bronchodilator) if wheezing
  - Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
  - If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
  - Alert emergency contacts.
  - Transport them to ER even if symptoms resolve. Person should remain in ER for at least 4 hours because symptoms may return.

**MILD SYMPTOMS**



**NOSE**  
Itchy/runny  
nose,  
sneezing



**MOUTH**  
Itchy mouth



**SKIN**  
A few hives,  
mild itch



**GUT**  
Mild nausea/  
discomfort

FOR MILD SYMPTOMS FROM MORE THAN ONE  
SYSTEM AREA, GIVE EPINEPHRINE.

FOR MILD SYMPTOMS FROM A SINGLE SYSTEM  
AREA, FOLLOW THE DIRECTIONS BELOW:

1. Antihistamines may be given, if ordered by a healthcare provider.
2. Stay with the person; alert emergency contacts.
3. Watch closely for changes. If symptoms worsen, give epinephrine.

**MEDICATIONS/DOSES**

Epinephrine Brand: \_\_\_\_\_

Epinephrine Dose: [ ] 0.15 mg IM [ ] 0.3 mg IM

Antihistamine Brand or Generic: \_\_\_\_\_

Antihistamine Dose: \_\_\_\_\_

Other (e.g., inhaler-bronchodilator if wheezing): \_\_\_\_\_

PARENT/GUARDIAN AUTHORIZATION SIGNATURE

DATE

PHYSICIAN/HCP AUTHORIZATION SIGNATURE


DATE

Please note: This is not a medication consent. A separate medication consent form is required.




### EPIPEN® (EPINEPHRINE) AUTO-INJECTOR DIRECTIONS

1. Remove the EpiPen Auto-Injector from the plastic carrying case.
2. Pull off the blue safety release cap.
3. Swing and firmly push orange tip against mid-outer thigh.
4. Hold for approximately 10 seconds.
5. Remove and massage the area for 10 seconds.




### AUVI-Q™ (EPINEPHRINE INJECTION, USP) DIRECTIONS

1. Remove the outer case of Auvi-Q. This will automatically activate the voice instructions.
2. Pull off red safety guard.
3. Place black end against mid-outer thigh.
4. Press firmly and hold for 5 seconds.
5. Remove from thigh.



### ADRENACLICK®/ADRENACLICK® GENERIC DIRECTIONS

1. Remove the outer case.
2. Remove grey caps labeled "1" and "2".
3. Place red rounded tip against mid-outer thigh.
4. Press down hard until needle penetrates.
5. Hold for 10 seconds. Remove from thigh.



### OTHER DIRECTIONS/INFORMATION (may self-carry epinephrine, may self-administer epinephrine, etc.):

1. Does this student need to sit at the nut-safe table in the lunchroom?  
 YES     NO
2. Does this student's classroom need to be nut-safe?  
 YES     NO

Treat the person before calling emergency contacts. The first signs of a reaction can be mild, but symptoms can get worse quickly.

### EMERGENCY CONTACTS — CALL 911

RESCUE SQUAD: \_\_\_\_\_

DOCTOR: \_\_\_\_\_ PHONE: \_\_\_\_\_

PARENT/GUARDIAN: \_\_\_\_\_ PHONE: \_\_\_\_\_

### OTHER EMERGENCY CONTACTS

NAME/RELATIONSHIP: \_\_\_\_\_

PHONE: \_\_\_\_\_

NAME/RELATIONSHIP: \_\_\_\_\_

PHONE: \_\_\_\_\_

PARENT/GUARDIAN AUTHORIZATION SIGNATURE

DATE

MUSKEGO-NORWAY SCHOOL DISTRICT  
Food Allergy/Anaphylaxis Emergency Action Plan

*\*Additional Information\**



Student's Name: \_\_\_\_\_

School: \_\_\_\_\_ Grade: \_\_\_\_\_

1. Please describe what reaction(s) this student has had in the past to the allergen(s) listed on the emergency action plan (itching, difficulty breathing, hives, etc.): \_\_\_\_\_

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

2. Has this student ever received an Epi Pen, Auvi-Q, or other source of epinephrine due to a severe allergic reaction? If yes, please describe the situation.

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

3. Any other special considerations for school:

\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_  
\_\_\_\_\_

4. (If you've not already addressed on the emergency action plan:)  
Does your student need to sit at the nut-safe table in the lunchroom?

Yes \_\_\_\_\_ No \_\_\_\_\_

Does your student's classroom need to be nut-safe?

Yes \_\_\_\_\_ No \_\_\_\_\_

Parent Signature: \_\_\_\_\_

Date: \_\_\_\_\_